

TURBULENCE IN CMEs

A. Ruzmaikin ; J. Feynman; W. J. Smith (all at: JPL, 4800 Oak Grove Dr., Pasadena, CA 91109; ph. 818-393-3953; e-mail: aruzmaikin@jplsp.jpl.nasa.gov)

Solar wind structures associated with coronal mass ejections have many distinguishing attributes. These attributes include for example, the relative abundance of helium, bidirectional electron streaming and high intensity helical largescale magnetic fields. Another distinguishing attribute may be the character of the MHD turbulence inside CMEs. We study fluctuations of the CME magnetic fields measured by ISHE3 and Ulysses spacecraft. The spectra and high-order structure functions characterizing the intermittency are found. The results are compared with those obtained for the quasi-stationary solar wind. The anisotropy of fluctuations in CMEs is estimated. This anisotropy is higher than that in the interplanetary medium due to the stronger magnetic fields. The helical properties of the turbulence inside CMEs are discussed.

American Geophysical Union Abstract Form

Reference # 0000
Session 0.00

1. 1996 Fall Meeting
2. 012406467
3. (a) Alexander Ruzmaikin
JPL, m/s 169-506,
4800 Oak Grove Dr.
Pasadena, CA 91106 (JSA
aruzmaikin@jplsp.jpl.nasa.gov
(b) 818-393-3953
(c) 1s1-351-8895
4. SH
5. (a) S1103
(1) 2131,2149,7.519
(c)
6. s / A
7. 0% published elsewhere
8. Charge \$50 to Alexander
Ruzmaikin MC card 5308 5172
0011 8850, expires 04/98
9. C
10. NONE
11. Regular author

Date received: 25 JUL 95
Date formatted: September 9, 1996
Form version: 1.3